

Test Description

The MolQ Germline Cancer Predisposition Panel include genes associated with both common and rare hereditary cancers.

Patient Demographic

Name: Mrs. Neena Walia
Sex: Female
Date of Birth/Age: 82 years
Disease: Pancreatic Carcinoma

Clinician

Clinician Name: Dr Amit Verma
Medical Facility: Dr AV Institute of Personalized Cancer Therapy and Research
Pathologist: Not Provided

Specimen

Booking ID: 012409130020
Sample Type: Blood
Tumor Content Percentage: NA
Date of Collection: 13-09-2024
Date of Booking: 13-09-2024

CLINICAL SYNOPSIS

Neena Walia, is a known case of pancreatic carcinoma and has a family history. Her sister had breast carcinoma. She has been evaluated for pathogenic variations in the genes listed in Appendix 2.

RESULTS

No pathogenic variant identified in relation to the phenotype.

PRIMARY FINDINGS

Gene (Transcript)#	Location	Variant	Zygosity	OMIM Phenotype	Inheritance	Clinical Significance [§]
No clinically relevant variant identified						

[§]Genetic test results are reported based on the recommendations of American College of Medical Genetics.

ADDITIONAL FINDINGS

Gene (Transcript)#	Location	Variant	Depth/ VAF	Zygosity	OMIM Phenotype	Inheritance	Clinical Significance [§]
G6PD (-) NM_001042351.3 (chrX:g.154533044C>T)	Exon 9	c.949G>A (p.Glu317Lys)	34X/ 47%	Heterozygous	G6PD deficiency	X-linked	Likely pathogenic (PM1, PP2, PP3, PS1)

[§]Genetic test results are reported based on the recommendations of American College of Medical Genetics.

G6PD DEFICIENCY

G6PD deficiency is the most common genetic cause of chronic and drug-, food-, or infection-induced hemolytic anemia. G6PD catalyzes the first reaction in the pentose phosphate pathway, which is the only NADPH-generation process in mature red cells; therefore, defense against oxidative damage is dependent on G6PD. Most G6PD-deficient patients are asymptomatic throughout their life, but G6PD deficiency can be life threatening. The most common clinical manifestations of G6PD deficiency are neonatal jaundice and acute hemolytic anemia, which in most patients is triggered by an exogenous agent, e.g., primaquine or fava beans. Acute hemolysis is characterized by fatigue, back pain, anemia, and jaundice. Increased unconjugated bilirubin, lactate dehydrogenase, and reticulocytosis are markers of the disorder. The striking similarity between the areas where G6PD

deficiency is common and Plasmodium falciparum malaria¹ is endemic provided evidence that *G6PD* deficiency confers resistance against malaria².

RECOMMENDATIONS

- Segregation analysis of the variant by Sanger sequencing is recommended in affected and unaffected members of the family.
- Validation of the variant(s) by Sanger sequencing is recommended to rule out false positives.
- Genetic counselling is advised for interpretation on the consequences of the variant(s).
- The significance/classification of the variants might change based on parental and family members genetic testing.
- For questions about this report, or for assistance in locating nearby genetic counseling services, please contact the laboratory: contact@molq.in.
- If results obtained do not match the clinical findings, additional testing should be considered as per referring clinician's recommendations.

REFERENCES

1. <https://omim.org/entry/611162>
2. Cappellini M D and Fiorelli G. Glucose-6-phosphate dehydrogenase deficiency. Lancet. 2008 Jan 5;371(9606):64-74.



Jatinder Kaur, PhD
Head, Molecular Biology & Genomics



Dr. Gulshan Yadav, MD
Head, Pathology

APPENDIX 1: TEST METHODOLOGY

Method

Next Generation Sequencing:¹⁻⁵ DNA extracted from blood, saliva, amniotic fluid, CVS or any other standard source is used for targeted capture-based Library preparation. Targeted capture provides an efficient and sensitive means for sequencing specific genomic regions in a high-throughput manner. The libraries were sequenced to mean >85-100x coverage on Illumina Novaseq 6000 sequencing platform with Paired End 2x150 chemistry.

Illumina DRAGEN Bio-IT Platform was followed for identification of variants in the sample. The sequences obtained are assembled and aligned to reference sequences based on NCBI RefSeq transcripts and human genome build (GRCh38). Haplotype caller has been used to identify variants which are relevant to the clinical indication.

In addition to SNVs and small Indels, copy number variants (CNVs) are detected from targeted sequence data using the commercially available algorithm. This algorithm detects rare CNVs based on comparison of the read-depths of the test data with the matched aggregate reference dataset. Clinically relevant mutations were annotated using published variants in literature, Commercial datasets and a set of diseases databases.

Common variants are filtered based on allele frequency in 1000Genome, ExAC, gnomAD, dbSNP and reference laboratory's internal database.

Non-synonymous variants effect is calculated using multiple algorithms such as PolyPhen-2, SIFT, MutationTaster2. Based on annotations data, ACMG rules-based classification performed for classification of variants identified through next generation sequencing study.

Average Sequencing Depth (x)	Percentage Target Base Pairs Covered		
	1x	≥5x	≥20x
86.41	99.00	98.80	98.44

Title	Data
Total data generated (Gb)	9.52
Total reads aligned (%)	100.00
Reads that passed alignment (%)	99.63
Data ≥ Q30 (%)	93.42

\$The classification of the variations is done based on American College of Medical Genetics as described below⁶

Variant	A change in a gene. This could be disease causing (pathogenic) or not disease causing (benign).
Benign	A variant which is known not to be responsible for disease has been detected. Generally, no further action is warranted on such variants when detected.
Likely Benign	A variant which is very unlikely to contribute to the development of disease however, the scientific evidence is currently insufficient to prove this conclusively. Additional evidence is expected to confirm this assertion of Pathogenicity.
Pathogenic	A disease-causing variation in a gene which can explain the patient's symptoms has been detected. This usually means that a suspected disorder for which testing had been requested has been confirmed.
Likely Pathogenic	A variant which is very likely to contribute to the development of disease however, the scientific evidence is currently insufficient to prove this conclusively. Additional evidence is expected to confirm this assertion of pathogenicity.
Variant of Uncertain Significance	A variant has been detected, but it is difficult to classify it as either pathogenic (disease causing) or benign (non-disease causing) based on current available scientific evidence. Further testing of the patient or family members as recommended by your clinician may be needed. It is probable that their significance can be assessed only with time, subject to availability of scientific evidence.

DISCLAIMER

- Interpretation of variants in this report is performed to the best knowledge of the laboratory based on the information available at the time of reporting. The classification of variants can change over time and MolQ cannot be held responsible

MolQ Laboratory (A Unit of Molecular Quest Healthcare Pvt. Ltd.)

Reference Laboratory: 28-29, Sector-18 (P) | Gurgaon, Haryana, 122015 | Phone 0124 - 4307906, Fax 0124 - 4278596 | Email: contact@molq.in

for this. Please feel free to contact MolQ Laboratory (contact@molq.in) in the future to determine if there have been any changes in the classification of any variations. Re-analysis of variants in previously issued reports in light of new evidence is not routinely performed, but may be considered upon request.

- The sensitivity of this assay to detect large deletions/duplications of more than 10 bp or copy number variations (CNV) is 70-75%. The CNVs detected with this assay have to be confirmed by alternate method such as MLPA & Microarray.
- Due to inherent technology limitations, coverage is not uniform across all regions. Hence pathogenic variants present in areas of insufficient coverage as well as those variants which currently do not correlate with the provided phenotype may not be analyzed/ reported. Additionally, it may not be possible to fully resolve certain details about variants, such as mosaicism, phasing, or mapping ambiguity.
- Genes with pseudogenes, paralog genes and genes with low complexity may have decreased sensitivity and specificity of variant detection and interpretation due to inability of the data and analysis tools to unambiguously determine the origin of the sequence data in such regions.
- The mutations have not been validated/confirmed by Sanger sequencing.
- Incidental or secondary findings (if any) that meet the ACMG guidelines⁷ can be given upon request.
- Sequence and copy number variants are reported according to the Human Genome Variation Society (HGVS).
- The transcript used for clinical reporting generally represents the canonical transcript, which is usually the longest coding transcript with strong/multiple supporting evidence. However, clinically relevant variants annotated in alternate complete coding transcripts could also be reported.
- The report shall be generated within turnaround time (TAT), however, such TAT may vary depending upon the complexity of test(s) requested. MolQ Laboratory under no circumstances will be liable for any delay beyond afore mentioned TAT.
- It is hereby clarified that the report(s) generated from the test(s) do not provide any diagnosis or opinion or recommends any cure in any manner. MolQ Laboratory hereby recommends the patient and/or the guardians of the patients, as the case may be, to take assistance of the clinician or a certified physician or doctor, to interpret the report(s) thus generated. MolQ Laboratory hereby disclaims all liability arising in connection with the report(s).
- In a very few cases genetic test may not show the correct results, e.g. because of the quality of the material provided to MolQ Laboratory. In case where any test provided by MolQ Laboratory fails for unforeseeable or unknown reasons that cannot be influenced by MolQ Laboratory in advance, MolQ Laboratory shall not be responsible for the incomplete, potentially misleading or even wrong result of any testing if such could not be recognized by MolQ Laboratory in advance.
- This is a laboratory developed test and the development and the performance characteristics of this test was determined by reference laboratory.

LIMITATIONS

- Genetic testing is an important part of the diagnostic process. However, genetic tests may not always give a definitive answer. In some cases, testing may not identify a genetic variant even though one exists. This may be due to limitations in current medical knowledge or testing technology. Accurate interpretation of test results may require knowing the true biological relationships in a family. Failing to accurately state the biological relationships in {my/my child's} family may result in incorrect interpretation of results, incorrect diagnoses, and/or inconclusive test results.
- Test results are interpreted in the context of clinical findings, family history and other laboratory data. Only variations in genes potentially related to the proband's medical condition are reported. Rare polymorphisms may lead to false negative or positive results. Misinterpretation of results may occur if the information provided is inaccurate or incomplete.
- Triplet repeat expansions, translocations, large deletion or duplications and chromosomal rearrangements events are currently not reliably detected by next generation sequencing.
- This assay is not meant to interrogate most promoter regions, deep intronic regions, or other regulatory elements, other gene rearrangements like inversion or translocation and does not detect single or multiexon deletions or duplications.
- Genetic testing is highly accurate. Rarely, inaccurate results may occur for various reasons. These reasons include, but are not limited to: mislabelled samples, inaccurate reporting of clinical/medical information, rare technical errors or unusual circumstances such as bone marrow transplantation, blood transfusion; or the presence of change(s) in such a small percentage of cells that may not be detectable by the test (mosaicism).
- It should be noted that this test does not sequence all bases in a human genome, not all variants have been identified or interpreted, and this report is limited only to variants with evidence for causing or contributing to disease/clinical details provided to MolQ Laboratory.
- Testing has been performed assuming that the sample received belongs to the above named individual and any stated relationships between individuals are accepted as true.
- The results should be interpreted in the context of the patient's medical evaluation, family history and racial/ethnic

MolQ Laboratory (A Unit of Molecular Quest Healthcare Pvt. Ltd.)

Reference Laboratory: 28-29, Sector-18 (P) | Gurgaon, Haryana, 122015 | Phone 0124 - 4307906, Fax 0124 - 4278596 | Email: contact@molq.in

background. Please note that variant classification and/or interpretation may change over time if more information available. Reinterpretation of multi gene next generation sequencing data is recommended on an annual basis and may be requested by a medical provider.

REFERENCES

1. Meyer L.R. et al., The UCSC Genome Browser database: extensions and updates 2013. *Nucleic Acids Res.*, 41(D1):D64-9, 2013.
2. McKenna, A., et al., The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.*, 20(9): 1297-303, 2010.
3. 1000 Genomes Project Consortium et al., A global reference for human genetic variation. *Nature*, 526(7571): 68-74, 2015.
4. Lek M. et al., Analysis of protein-coding genetic variation in 60,706 humans. *Nature*, 536(7616):285-91, 2016.
5. McLaren W. et al., Deriving the consequences of genomic variants with the Ensembl API and SNP Effect Predictor. *Bioinformatics.*, 26(16):2069-70, 2010.
6. Richards S. *et al.*, Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology, *Genet Med.*, 17(5):405-24, 2015.
7. Kalia S.S. et al., Recommendations for reporting of secondary findings in clinical exome and genome sequencing, 2016 update (ACMG SF v2.0): a policy statement of the American College of Medical Genetics and Genomics. *Genet Med.*, 19(2):249-255, 2017.

APPENDIX 2: GENE LIST WITH COVERAGE

Gene	Percentage of coding region covered	Gene	Percentage of coding region covered	Gene	Percentage of coding region covered
ABRAXAS1	100.00	FANCI	100.00	PRSS1	100.00
ACD	100.00	FANCL	100.00	PTCH1	100.00
ACVRL1	100.00	FANCM	100.00	PTCH2	100.00
ADA	100.00	FAS	100.00	PTEN	100.00
AIP	100.00	FASLG	100.00	PTPN11	100.00
AKT1	100.00	FCHO1	100.00	RAC2	100.00
ALK	100.00	FGFR2	100.00	RAD50	100.00
ANKRD26	100.00	FH	100.00	RAD51	100.00
APC	100.00	FLCN	100.00	RAD51C	100.00
APCS	100.00	FOXO1	100.00	RAD51D	100.00
AR	100.00	G6PD	100.00	RAF1	100.00
ATM	100.00	GALNT12	100.00	RASA2	100.00
ATR	100.00	GATA2	100.00	RASAL1	100.00
AXIN1	100.00	GNA13	100.00	RASGRP1	100.00
AXIN2	100.00	GPC3	100.00	RB1	100.00
B2M	100.00	GPR101	100.00	RECQL	100.00
BAP1	100.00	GREM1	100.00	RECQL4	100.00
BARD1	100.00	HAVCR2	100.00	REST	100.00
BCL2	100.00	HNF1A	100.00	RET	100.00
BLM	100.00	HNF1B	100.00	RFWD3	100.00
BMPR1A	100.00	HOXB13	100.00	RHBDF2	100.00
BRAF	100.00	HRAS	100.00	RHOH	100.00
BRCA1	100.00	IKZF1	100.00	RINT1	100.00
BRCA2	100.00	IL2RA	100.00	RIT1	100.00
BRIP1	100.00	IL2RB	100.00	RMRP	0.00
BTNL2	100.00	ITK	100.00	RNASEL	100.00
BUB1B	100.00	KIF1B	100.00	RNF43	100.00
CARD11	100.00	KIT	100.00	RPS20	100.00
CARMIL2	100.00	KITLG	100.00	RRAS	100.00
CASP10	100.00	KLHDC8B	100.00	RUNX1	100.00
CASP8	100.00	KMT2D	100.00	SAMD9	100.00
CASR	100.00	KRAS	100.00	SAMD9L	100.00
CBL	100.00	LIG4	100.00	SBDS	100.00
CD27	100.00	LSP1	79.12	SDHA	100.00
CD70	100.00	LZTR1	100.00	SDHAF2	100.00
CD79A	100.00	MAD2L2	100.00	SDHB	100.00
CD79B	100.00	MAGT1	100.00	SDHC	100.00
CD82	100.00	MAP2K1	100.00	SDHD	100.00
CDC73	100.00	MAP2K2	100.00	SH2D1A	100.00
CDH1	100.00	MAP3K1	100.00	SHOC2	100.00
CDK4	100.00	MAX	100.00	SLX4	100.00
CDKN1B	100.00	MC1R	100.00	SMAD4	100.00
CDKN1C	100.00	MCM4	100.00	SMARCA2	100.00
CDKN2A	100.00	MEN1	100.00	SMARCA4	100.00
CDKN2B	100.00	MET	100.00	SMARCB1	100.00
CEBPA	100.00	MITF	100.00	SMARCE1	100.00

MolQ Laboratory (A Unit of Molecular Quest Healthcare Pvt. Ltd.)

Reference Laboratory: 28-29, Sector-18 (P) | Gurgaon, Haryana, 122015 | Phone 0124 - 4307906, Fax 0124 - 4278596 | Email: contact@molq.in

CEP57	100.00	MLH1	100.00	SOCS1	100.00
CFTR	100.00	MLH3	100.00	SOS1	100.00
CHEK2	100.00	MRE11	100.00	SOS2	100.00
CPA1	100.00	MSH2	100.00	SPINK1	100.00
CREBBP	100.00	MSH3	100.00	SPRED1	100.00
CTLA4	100.00	MSH6	100.00	SRP72	96.83
CTNNA1	100.00	MSR1	100.00	STAT3	100.00
CTPS1	100.00	MUTYH	100.00	STAT6	100.00
CYLD	100.00	MXI1	100.00	STK11	100.00
DDB2	100.00	MYC	100.00	STK4	100.00
DDX41	100.00	MYD88	100.00	STXBP2	100.00
DICER1	100.00	MYSM1	100.00	SUFU	100.00
DIS3L2	100.00	NBN	100.00	TERC	0.00
DKC1	100.00	NF1	100.00	TERF2IP	100.00
DOCK8	100.00	NF2	100.00	TERT	100.00
EFL1	100.00	NHEJ1	100.00	TET2	100.00
EGFR	100.00	NOTCH1	100.00	TGFB1	100.00
ELAC2	100.00	NOTCH2	100.00	TGFBR2	100.00
ELANE	100.00	NPAT	100.00	TINF2	100.00
ENG	100.00	NRAS	100.00	TMEM127	100.00
EP300	100.00	NSD1	100.00	TNFAIP3	100.00
EPCAM	100.00	NSUN2	100.00	TNFRSF13B	100.00
ERCC1	100.00	NTHL1	100.00	TNFRSF14	100.00
ERCC2	100.00	PALB2	100.00	TOX3	96.65
ERCC3	100.00	PALLD	100.00	TP53	100.00
ERCC4	100.00	PAX5	100.00	TPP2	100.00
ERCC5	100.00	PDGFRA	100.00	TRIP13	100.00
ETV6	100.00	PHOX2B	100.00	TSC1	100.00
EXO1	100.00	PIK3CA	100.00	TSC2	100.00
EXT1	100.00	PIK3CD	100.00	UBE2T	100.00
EXT2	100.00	PIK3R1	100.00	VHL	100.00
EZH2	100.00	PIM1	100.00	VPS13B	100.00
FADD	100.00	PMS1	100.00	WAS	100.00
FAM111B	100.00	PMS2	100.00	WRN	100.00
FAN1	100.00	POLD1	100.00	WT1	100.00
FANCA	100.00	POLE	100.00	XIAP	100.00
FANCB	100.00	POLH	100.00	XPA	100.00
FANCC	100.00	POT1	100.00	XPC	100.00
FANCD2	100.00	PPM1D	100.00	XRCC2	100.00
FANCE	100.00	PRF1	100.00	XRCC3	100.00
FANCF	100.00	PRKAR1A	100.00	ZFHX3	100.00
FANCG	100.00	PRKCD	100.00		

MTDNA GENES LIST WITH COVERAGE

Gene	Percentage of coding region covered	Gene	Percentage of coding region covered	Gene	Percentage of coding region covered
MT-ATP6	100.00	MT-ND6	100.00	MT-TL1	100.00
MT-ATP8	100.00	MT-RNR1	100.00	MT-TL2	100.00

MolQ Laboratory (A Unit of Molecular Quest Healthcare Pvt. Ltd.)

Reference Laboratory: 28-29, Sector-18 (P) | Gurgaon, Haryana, 122015 | Phone 0124 - 4307906, Fax 0124 - 4278596 | Email: contact@molq.in

<i>MT-CO1</i>	100.00	<i>MT-RNR2</i>	100.00	<i>MT-TM</i>	100.00
<i>MT-CO2</i>	100.00	<i>MT-TA</i>	100.00	<i>MT-TN</i>	100.00
<i>MT-CO3</i>	100.00	<i>MT-TC</i>	100.00	<i>MT-TP</i>	100.00
<i>MT-CYB</i>	100.00	<i>MT-TD</i>	100.00	<i>MT-TQ</i>	100.00
<i>MT-ND1</i>	100.00	<i>MT-TE</i>	100.00	<i>MT-TR</i>	100.00
<i>MT-ND2</i>	100.00	<i>MT-TF</i>	100.00	<i>MT-TS1</i>	100.00
<i>MT-ND3</i>	100.00	<i>MT-TG</i>	100.00	<i>MT-TS2</i>	100.00
<i>MT-ND4</i>	100.00	<i>MT-TH</i>	100.00	<i>MT-TT</i>	100.00
<i>MT-ND4L</i>	100.00	<i>MT-TI</i>	100.00	<i>MT-TV</i>	100.00
<i>MT-ND5</i>	100.00	<i>MT-TK</i>	100.00	<i>MT-TW</i>	100.00
<i>MT-TY</i>	100.00				